MBMG
Metabarcoding & Metagenomics

Metabarcoding and Metagenomics 10: 163–189 (2026)
DOI: 10.3897/mbmg.10.174947

Research Article

DNA barcoding reference dataset revealed overlooked diversity and clarified biogeography of fishes in the largest European river system – Volga River

Boris A. Levin[1], Marina A. Levina[1], Oleg N. Artaev[2], Aleksey A. Bolotovskiy[2], Alexander A. Gandlin[1], Ilya S. Turbanov[1,2,3], Aleksandra S. Komarova[1,2], Nikolai S. Mugue[4], Ivan V. Pozdeev[5], Alexander B. Ruchin[6], Kirill V. Litvinov[7], Stepan A. Podolyako[8], Evgeniy Simonov[1]

1 *A.N. Severtsov Institute of Ecology and Evolution of the Russian Academy of Sciences, Moscow, Russia*
2 *Papanin Institute for Biology of Inland Waters, Russian Academy of Sciences, Borok, Russia*
3 *T.I. Vyazemsky Karadag Scientific Station, Nature Reserve of the RAS, Branch of A.O. Kovalevsky Institute of Biology of the Southern Seas of RAS, Theodosia, Crimea*
4 *Russian Federal Research Institute of Fisheries and Oceanography, Moscow, Russia*
5 *Russian Federal Research Institute of Fisheries and Oceanography, (Perm Branch), Perm, Russia & Saint-Petersburg Research Center of the Russian Academy of Sciences, Saint-Petersburg, Russia*
6 *Mordovia State Nature Reserve and National Park «Smolny», Republic of Mordovia, Saransk, Russia*
7 *Astrakhan State Reserve, Astrakhan, Russia*
8 *Shirshov Institute of Oceanology, Russian Academy of Science, Moscow, Russia*

Corresponding author: Boris A. Levin (borislyovin@gmail.com)

Abstract

This study represents the first thorough genetic assessment of freshwater fishes from the Volga River Basin – the longest (3,690 km) river in Europe. DNA barcode sequences (*COI*) were studied in 84 species (approx. 85% of all diversity) belonging to 57 genera and 25 families. In total, 1,327 DNA barcodes were used to: (i) build a DNA barcode reference library and check for the presence of a barcoding gap; ii) assess the genetic and taxonomic diversity of Volga fishes and iii) clarify the biogeography of Volga fishes. DNA barcodes are reliable means for the identification of over 95% of analysed species showing barcode gaps. An overlap between intra- and interspecific genetic distances was present in four species of genera *Leuciscus* and *Rutilus*. DNA barcoding identified seven new species for the Volga. Amongst these, two are aggressive alien species (*Hemiculter leucisculus* and *Pseudorasbora parva*), while five are indigenous to the region and belong to the genera *Salmo*, *Leuciscus*, *Gobio* and *Cobitis* from neighbouring basins, which naturally or human-aided colonised the Volga Basin. Moreover, three new phylogenetic lineages were discovered in the genera *Leucaspius*, *Phoxinus* and *Proterorhinus*. While the fishes from sub-basins of the Volga usually do not demonstrate sorting of the haplotypes, the more ancient Kama displays regional endemism for rheophilic species (e.g. *Barbatula*, *Cottus*). Our results suggest the Kama might serve as refuge during Pleistocene climate oscillations. In summary, the generated *COI* dataset fills the existing gap in *COI* database of fishes inhabiting freshwaters of Eastern Europe.

Key words: Cryptic diversity, cytochrome c oxidase subunit I, Eastern Europe, mitochondrial DNA, molecular identification, phylogeography

OPEN ACCESS

Academic editor: Florian Leese
Received: 15 October 2025
Accepted: 12 February 2026
Published: 23 February 2026

Citation: Levin BA, Levina MA, Artaev ON, Bolotovskiy AA, Gandlin AA, Turbanov IS, Komarova AS, Mugue NS, Pozdeev IV, Ruchin AB, Litvinov KV, Podolyako SA, Simonov E (2026) DNA barcoding reference dataset revealed overlooked diversity and clarified biogeography of fishes in the largest European river system – Volga River. Metabarcoding and Metagenomics 10: e174947. https://doi.org/10.3897/mbmg.10.174947

Copyright: © Boris A. Levin et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Biodiversity is a cornerstone for the sustainable functioning of ecosystems (Bengtsson et al. 2000; Hooper et al. 2005; Cardinale et al. 2012). Fishes are the most diverse group of vertebrates with a species number exceeding 37,000 (Fricke et al. 2025), although their diversity is still significantly underestimated (e.g. Zemlak et al. (2009); Wang et al. (2020); Tsoupas et al. (2022)). Despite freshwater ecosystems cover only 0.8% of the Earth's surface, they disproportionately contribute to approximately half of the global known fish diversity (Barbarossa et al. 2021). In addition, fishes are an important food source that, in turn, need proper management for sustained resource use. Freshwater habitats are more threatened by human activities and environmental change, being more vulnerable compared to those in marine or terrestrial ecosystems (Sayer et al. 2025). Freshwater ecosystems in densely populated areas are degraded faster than in less-altered environments. The human pressure on aquatic ecosystems can result in the extinction of a significant part of fish diversity before it is scientifically documented (Dodds et al. 2013).

DNA barcoding is a well-established tool for species identification, biodiversity assessment and monitoring of ecosystems, significantly contributing to the understanding of fish species diversity (Hebert and Gregory 2005; Ward et al. 2009; Knebelsberger et al. 2015; Behrens-Chapuis et al. 2021; Sheraliev and Peng 2021; Tsoupas et al. 2022; Zangl et al. 2022; Appleyard et al. 2025). It is beneficial compared to morphology-based identification especially when cryptic/pseudocryptic species, larval or juvenile stage fish or parts of organisms are analysed (Steinke et al. 2009). In addition, data on genetic polymorphism of broadly distributed species, based on DNA barcodes, may provide information on population structure, genetic diversity, evolutionary history and might be helpful in the delineation of conservation units. Such analyses are, in particular, supported through novel and non-invasive environmental DNA (eDNA) based approaches (Ushio et al. 2018; Snyder et al. 2020; Zhang et al. 2024). However, the accuracy of eDNA approaches may be greatly restricted by some factors, amongst which an absence or insufficiency of a comprehensive DNA barcode reference database (Marques et al. 2021; Xiong et al. 2022; Zhang et al. 2025).

The Volga is the longest European river (3,690 km) with a catchment area of about 1,360,000 km^2. Volga flows through heterogeneous ecoregions from north to south – forests, forest steppe, steppe and semi-desert and borders the Azov and Black Sea Basin, Baltic Sea Basin, White Sea Basin, Ob' River system (Kara Sea) and the Ural River system in the Caspian Sea drainage (Mineeva et al. 2022). Water ecosystems in the Volga drainage are severely threatened by anthropogenic pressure because of their location in an economically developed and densely populated area - ca. 70 million people dwell in the Volga Basin (Mineeva et al. 2022). In addition, the construction of numerous dams on the Volga River in the 20th century drastically altered its hydrological regime and interrupted or shortened the migration routes of anadromous fish, including valuable species of sturgeons (Ruban et al. 2019). All of this makes the indigenous fish fauna of the Volga River highly vulnerable to various direct and indirect human impacts, including the influence of alien species (Panov et al. 2007; Schletterer et al. 2018; Mineeva et al. 2022).

Unexpectedly, fish diversity estimates for the Volga Basin largely vary in modern literature – from 79 (Schletterer et al. 2018) and 96 (Dyldin et al. 2020) to 114 (Slynko and Tereshchenko 2014) and even 117 (Froese and Pauly 2025).

In our view, the two latter counts are greatly overestimated since they include all cases of historical introductions, naturalisation which have never been reported (e.g. *Oncorhynchus* spp., *C. baunti*, *Parabramis pekinensis* etc.). They also list the species from the surrounding waters in the Caspian Sea, but never recorded in the Volga (e.g. *Acipenser persicus*, *Gasterosteus aculeatus* - Froese and Pauly (2025)) or mistakenly listed species (e.g. *Proterorhinus marmoratus* that was listed along with *P. nasalis*; *Ameiurus nebulosus* - Froese and Pauly (2025)) or listed species of uncertain, probably, hybrid origin - for example, *Alosa suworowi* (see Dyldin et al. (2020)). According to our preliminary assessment, based on the literature survey and own sampling during the last 25 years, the list of native and non-native fishes and lampreys inhabiting the Volga Basin includes ca. 90 species, of which 78 are indigenous and 12 are considered alien.

Despite the high interest in Volga's fish diversity, only few genetic studies have been devoted to the indigenous Volga fishes (Mendel et al. 2008; Levin et al. 2016, 2017, 2018; Borovikova 2017; Artaev et al. 2021, 2024). For example, only 179 sequences of DNA barcodes of *COI* have been deposited in NCBI's GenBank so far (30 Sept 2025), 108 of which are from invasive fish fauna (Karabanov et al. 2022). The lack of DNA barcodes of Volga's fishes strongly impedes the implementation of various genetic approaches to identify species, for example, via eDNA, for various purposes – monitoring of biodiversity, surveys of endangered or alien species (including eDNA applications), food control and forensic research.

In view of these limitations, the goal of our study was fourfold: i) to close the existing gap in the COI DNA barcode reference library for the majority of Volga fish species; ii) to characterise mitochondrial genetic diversity; iii) to update the taxonomy of Volga species and iv) to clarify the biogeography of Volga fishes, based on new knowledge obtained.

Materials and methods

Sampling

Fishes were sampled during field studies in 2011-2024. In particular, 1151 specimens from 80 species were collected from 202 sampling sites in all Volga sub-basins according to Mineeva et al. (2022): i) Upper Volga; ii) Middle Volga; iii) Kama and iv) Lower Volga (Fig. 1. Suppl. material 1: table S1). Fin clips or scales were taken and stored in 96% ethanol. For most fin-clips, voucher specimens were stored at the Fish Collection of the Papanin Institute for Biology of Inland Waters of the Russian Academy of Sciences (**IBIW_FS**), Borok, Russia. DNA samples of protected species (except for some *Acipenser ruthenus*) were taken from the collection of the Russian Federal Research Institute of Fisheries and Oceanography, Moscow, Russia.

Specimens were identified in the field or in the lab, based on morphological traits using the most recent identification keys or species descriptions (Kottelat and Freyhof 2007; Mendel et al. 2008; Neilson, Stepien 2009a; Mousavi-Sabet et al. 2011; Bogutskaya et al. 2013; Levin et al. 2016, 2017, 2018; Pereira et al. 2021; Segherloo et al. 2021; Artaev et al. 2024; Turbanov et al. 2025; Vasil'eva et al. 2025). Examined species include nine species protected by the IUCN Red List of Threatened Species (2025) and the Red Book of Russian Federation (2021): *Acipenser ruthenus* (EN), *A. gueldenstaedtii* (CR), *A. nudiventris* (CR), *A. stellatus* (CR), *Huso huso* (CR), *Coregonus albula* (EN - population of Lake Pleshchejevo

Figure 1. Map of fish sampling localities. Roman numbers designate four Volga's sub-basins: I – Upper Volga, II – Middle Volga, III – Kama and IV – Lower Volga. Arabic numbers display sampling localities; geographic coordinates and other details of sampling localities are given in Suppl. material 1: table S1.

in the Volga Basin is protected by the national Red Book), *Stenodus leucichthys* (NT), *Hucho taimen* (VU) and *Salmo caspius* (CR - resident populations in the Middle Volga and Kama are protected by the national Red Book).

DNA extraction, PCR and sequencing

DNA was extracted from 1151 samples belonging to 56 genera and 80 species, using the salt-extraction protocol of Aljanabi and Martinez (1997). The numbering of the samples, the species to which they belong and their sampling sites are

given in Suppl. material 1: table S1. The barcoding fragment of mitochondrial *COI* gene was amplified with Polymerase Chain Reactions (PCR), using an M13-tailed primer cocktail with the primers VF2_t1, FishF2_t1, FishR2_t1 and FR1d_t1 (Ivanova et al. 2007) or LCO 1490 and HCO 1298 primers (Folmer et al. 1994) for some fish taxa; lampreys were amplified and sequenced using the primers COI-F1 (Ivanova et al. 2007) and COI-2R (Ward et al. 2005). A standard 25 µl PCR mix consisted of 2.5 µl PCR buffer (10x); 2.5 µl dNTP (2 mM); 1.5 µl 'Fish Cocktail' (2 µM); 0.2 Taq DNA Polymerase (5 units per µl); 16.75 µl mQ-H2O and 1.0 µl of the extracted DNA sample. The temperature profile for various amplification protocols is given in Suppl. material 1: table S2. PCR products were visually checked on 1.5% agarose gels and purified using 96% ethanol and sodium acetate (3 M). Sanger sequencing using M13 primers, LCO1490 and HCO1298 primers, as well as COI-F1 and COI-2R primers, was performed in IBIW RAS using an ABI's 3500 DNA Analyser. All sequences were deposited to GenBank under Accession numbers PV840619–PV841769 (Suppl. material 1: table S1).

Barcoding analyses

Newlyobtained barcode sequences along with the data available in GenBank were combined, totalling of 1327 individuals. Of these, 176 *COI* sequences were retrieved from GenBank (Neilson and Stepien 2009a, 2009b; April et al. 2011; Haponski and Stepien 2013; Behrens-Chapuis et al. 2015; Kodukhova et al. 2016; Levin et al. 2016; Mousavi-Sabet et al. 2016; Levin et al. 2018; Palandačić et al. 2020; Kodukhova and Karabanov 2021; Weiss et al. 2021; Karabanov et al. 2022; Karabanov et al. 2023; Artaev et al. 2024; Turbanov et al. 2025, as well as unpublished sequences deposited to NCBI; see Suppl. material 1: table S1 for details). Multiple sequence alignment (MSA) was created with MUSCLE 3.8.31 (Edgar 2004). All sequences were then translated to protein and checked for the presence of unexpected stop codons. To examine the clustering of DNA barcode sequences in a gene tree, we used the Maximum Likelihood (ML) approach in IQ-TREE 2.4.0 (Minh et al. 2020). In total, 312 unique haplotypes were used. The best-fit models and partitioning scheme by codon position were selected with ModelFinder (Chernomor et al. 2016; Kalyaanamoorthy et al. 2017) using Bayesian Information Criterion (BIC) as follows: TIM3+F+I+G4 for codon position 1, F81+F+R2 for codon position 2 and TN+F+R4 for codon position 3. Tree support was accessed with 1 000 ultrafast bootstrap replicates (Hoang et al. 2018) and 1000 replicates of the Shimodaira–Hasegawa-like approximate likelihood ratio test (SH-aLRT; Guindon et al. (2010)).

To examine the species identification success, we applied Best match (BM), Best close match (BCM) and All Species Barcodes (ASB) approaches (Meier et al. 2006) as implemented in SpeciesIdentifier v. 1.9 (http://taxondna.sourceforge.net). The Automatic Barcode Gap Discovery (ABGD) method (Puillandre et al. 2012) for the detection of the "barcoding gap" in the distribution of Kimura 2-parameter (K2P) pairwise distances in the dataset was used. The ABGD was run with the default parameters, except for K2P distance (option -d 0). In addition, we used the recently developed Assemble Species by Automatic Partitioning (ASAP) method (Puillandre et al. 2012), which relies on a hierarchical clustering algorithm of pairwise genetic distances to build species partitions. The ASAP was run on the web server (https://bioinfo.mnhn.fr/abi/public/asap/asapweb.html) with split groups below 0.05 probability and K2P distances. Additionally,

we checked the dataset for the presence of the "barcoding gap" (Meyer and Paulay 2005) by calculating the furthest intraspecific distance and the closest non-conspecific distance ("nearest neighbour") for each individual in the dataset using Spider v. 1.5.1 R-package (Brown et al. 2012).

To show different phylogeny patterns, a haplotype network was constructed using the median joining algorithm (Bandelt et al. 1999) in POPART 1.7 (Leigh and Bryant 2015) with the default value of epsilon (0).

Results

A total of 1151 specimens from the Volga system were successfully newly barcoded in this study. They belonged to 25 families, 56 genera and 80 species. The comprehensive dataset consisted of our and previously published data totalling 1327 individuals of 57 genera and 84 species (Table 1, Fig. 2, Suppl. material 3). After alignment and trimming, all barcode sequences were 643 bp long without any ambiguities or missing sites. The mean nucleotide frequencies of the entire dataset were A (24.1%), T (29.1%), G (19.1%) and C (27.8%). The genetic distance K2P within species ranged from 0.0 to 2.4%.

For 14 species, *Atherina caspia*, *Babka gymnotrachelus*, *Benthophilus granulosus*, *Caspiosoma caspium*, *Gobio caucasicus*, *Hypophthalmichthys molitrix*, *Knipowitschia longecaudata*, *Leuciscus baicalensis*, *Neogobius pallasi*, *Phoxinus* sp., *Ponticola syrman*, *Pseudorasbora parva*, *Pungitius pungitius* and *Vimba vimba*, only a single sequence was obtained. Of all other species, at least two, but mostly substantially more sequences (3–150, average 15.8) were obtained, providing the opportunity to assess intraspecific variation (Table 1, Suppl. material 1: table S).

Barcoding analyses

Identification success rates using BM, BCM and ABS approaches at 2% K2P threshold distance are provided in Table 2. The ASB approach was less powerful with numerous ambiguous identifications (102 out of 1327, 7.68%). A small fraction (16 out of 1327, 1.2%) of ambiguous matches in BM and BCM were in *Rutilus caspius* / *R. lacustris* and *Leuciscus idus* / *L. leuciscus*. All incorrect identifications in BM and absence of any matches closer than the threshold in BCM and ABS were due to the singleton species in the dataset. In BCM and ABS analyses, a single incorrect identification was for *Knipowitschia longecaudata* (identified as *K. caucasica* with 1.09% distance).

Table 1. Summary of basic barcode statistics.

Statistics	Value
Total individuals	1327
This study	1151
Previously published	176
Number of species	84
Number of genera	57
Mean number of individuals per species (range)	15.8 (1-150)
Number of singletons	14
Mean intraspecific K2P distance (range)	0.21% (0.0-2.38%)
Mean interspecific, intrageneric K2P distance (range)	5.69% (0.0-15.77%)



Figure 2. Pie chart showing the taxonomic composition of the examined *COI* dataset of fishes from the Volga River Basin by family, genus and species. Interactive version of the plot is available as a Suppl. material 2.

Table 2. Results of identification success analysis using various criteria. BM – best match, BCM – best close match, ASB – all species barcodes; threshold is 2% K2P distance.

Identification result	BM, %	BCM, %	ASB, %
Correct	97.73	97.73	91.25
Ambiguous	1.2	1.2	7.68
Incorrect	1.05	0.07	0.07
Sequences without any match closer than threshold	NA	0.97	0.97

ASAP yielded 84 subsets/species with the lowest (best) ASAP score of 7.0, at the threshold distance of 0.035 (see Suppl. material 1: table S5). The identical subsets were also produced in a range of threshold distances from 0.015 to 0.032. While this number (84) corresponds to the total number of predefined species in the specified dataset, it does not exhibit complete concordance due to the presence of both split and lumped taxa. *Benthophilus* species are lumped into one group, as well as in *Knipowitschia*, *Leuciscus idus / L. leuciscus* and *Rutilus caspicus / R. lacustris*. On the other hand, *Carassius gibelio*, *Leucaspius delineatus*, *Perccottus glenii* and *Squalius cephalus* species each split into two groups. The ABGD recursive partitions with prior maximal distance (P) of 0.013, 0.022 returned 80 groups, while a recursive partition with P = 0.008 showed 85 groups, which is close to the number of predefined species in the dataset. However, the resulting partitions also split or combined some predefined species (see Suppl. material 1: table S6). For instance, with 80 groups, the species

in *Benthophilus*, *Knipowitschia*, *Leuciscus idus / L. leuciscus*, *Rutilus caspicus / R. lacustris* and *Salmo* are lumped, while *Leucaspius delineatus* split into two groups (see also Figs 3, 4).

Phylogeography patterns

Various phylogeography patterns of Volga's fishes were revealed. Below, we consider four of the most representative examples of phylogeography patterns (Fig. 5). A single haplotype was for the limnophilous species - rudd *Scardinius erythrophthalmus* (n = 22) in spite of its broad geographic distribution. In particular, studied sites are ranged from the Upper Volga (58.4754 N, 35.4646 E) to the Lower Volga (46.2752 N, 47.8622 E) (Fig. 5A). Contrarily, highly polymorphic, but geographically not structured haplotype net is for the rheophilic species, common dace *Leuciscus leuciscus* (n = 78; 21 haplotypes). Although the haplotype net of



Figure 3. DNA barcode (*COI*) gene tree inferred using the Maximum Likelihood method. The tree comprises 312 unique haplotypes from 84 species of fish and lampreys from the Volga River Basin. Numbers near the nodes are SH-aLRT support (%) / ultrafast bootstrap support (%) above 50%. The units of the scale bar and branch lengths are expected substitutions per site. Red colour is for species (or new evolutionary lineages, possibly cryptic species) detected in the Volga Basin for the first time.



Figure 4. Barcode gap plot of the relationship between the distance to the closest non-conspecific (nearest neighbour) and maximum intraspecific K2P distance calculated as an average for each species with sample size above one. Dots above the 1:1 red line indicate the (likely) presence of a barcode gap, i.e. greater interspecific than intraspecific divergences.



Figure 5. Haplotype networks of four selected species of the Volga Basin, represented different phylogeographical patterns. **A.** Monomorphic *Scardinius erythrophthalmus* (Leuciscidae); **B.** Diversified, but not geographically structured of *Leuciscus leuciscus* (Leuciscidae); **C.** Diversified and partially geographically structured of *Barbatula barbatula* (Nemacheilidae); **D.** Not diversified, but strictly geographically subdivided of *Cottus koshewnikowi* (Cottidae). Black dots represent hypothetical intermediate haplotypes. Each short line signifies a single nucleotide substitution.

L. leuciscus is complex, fish from the Upper and Middle Volga as well as from the Kama basins share the same haplotypes (Fig. 5B). Another phylogeography pattern – similar to that of *L. leuciscus* in terms of haplotype diversity, but showing some geographical subdivision within the Volga Basin – is for more rheophilic

species preferring the clear and cold water, the stone loach *Barbatula barbatula* (n = 64; 18 haplotypes – Fig. 5C). In particular, most of the Kama samples are sorted mainly in one of two haplogroups, while samples from the Upper and Middle Volga share both haplogroups (Lower Volga is represented by a few samples). Another pattern of phylogeography - lower haplotype diversity, but strong sorting in geography – is displayed by even more rheophilic and more cold-water species, the Koshewnikow's sculpin *Cottus koshewnikowi* (n = 40; 6 haplotypes – Fig. 5D). Its Kama's populations are moderately diverse (three haplotypes) as populations from the Upper and Middle Volga taken together (*Cottus* is absent in the Lower Volga), but Kama and Volga samples do not share the haplotypes (Fig. 5D). Moreover, some populations of the Kama are rather divergent from Volga populations (six mutational steps) that can be considered as species level divergence for European sculpins of the genus *Cottus*, characterised by low interspecific divergence levels in *COI* (Knebelsberger et al. 2015; Sideleva and Zhidkov 2024).

Discussion

Our study provided the first detailed and comprehensive genetic assessment of fishes and lampreys inhabiting the largest European riverine basin – the Volga River system. In total, we provided *COI* barcodes for approximately 85% of the species recorded in the Volga Basin. Some rare or episodically occurring species in the Lower Volga or Volga's estuary have not yet been studied. Amongst these are rare anadromous species [e.g. *Caspiomyzon wagneri* (Kessler, 1870), *Luciobarbus capito* (Güldenstädt, 1773)] or species inhabiting the brackish waters of the Caspian Sea and episodically recording in the estuary (e.g. *Alosa* spp., *Benthophilus* spp. and *Syngnathus caspius* Eichwald, 1831). Generated *COI* barcodes showed high identification success (> 95% of analysed species). This is a very high ratio given that the success rate of DNA barcoding in most studies ranges from 76.4% to 100% (e.g. Hubert et al. (2008); Chen et al. (2015); Sheraliev and Peng (2021); Tsoupas et al. (2022); Muhala et al. (2024)) and is comparable to that in the study of Zangl et al. (2022) for fish diversity in Austria, Central Europe. Therefore, obtained reference sequences might be used for various DNA-based techniques for purposes of species identification, biodiversity assessment and monitoring, phylogeography, phylogeny, population genetics studies, as well as conservation genetics and food control. In addition to creating a reference database, our study largely contributed to knowledge on the diversity of fishes in various aspects and clarified the biogeography of Volga fishes. We consider these issues in more detail below.

Novel fish species for the Volga Basin

Seven new species for the Volga Basin and three new phylogenetic lineages (candidates for new species) from six families were discovered in the Volga system: *Salmo trutta*, *Gobio caucasicus*, *Hemiculter leucisculus*, *Leucaspius* sp., *Leuciscus baicalensis*, *Phoxinus* sp., *Pseudorasbora parva*, *Cobitis faridpaki*, *Cobitis tanaitica* and *Proterorhinus* sp. Some of these were identified by morphology and confirmed by *COI* barcodes (e.g. *Hemiculter leucisculus*, *Pseudorasbora parva*), while some others were detected primarily by *COI* barcodes (cryptic or pseudocryptic diversity).

New alien species

Hemiculter leucisculus (n = 2; see photograph in Suppl. material 3) was recorded on 23 June 2021 in the Lower Volga (Lagan channel – locality no. 201 in Fig. 1). It is considered one of the very dangerous alien species that quickly establish populations in new areas and may affect local ecosystems (Dong et al. 2020). Previously established populations were detected in the neighbouring water systems in the Caspian Sea Basin in Azerbaijan, as well as the Daghestan Region of Russia (Zardoya and Doadrio 1999; Mustafayev et al. 2015). Novel finding suggests an expansion of *H. leucisculus* in the system of Caspian Sea. Further monitoring of the expansion of this alien species is highly required in the Lower Volga. *Pseudorasbora parva* is another aggressive alien species that is quickly colonising the Volga Basin. One individual (see photograph in Suppl. material 3) was recorded on 25 August 2024 in the Vershaut River, Middle Volga (locality no. 151 in Fig. 1), which shares watershed with the Don Basin tributaries, where this species was known from the previous study (Karabanov et al. 2009). Another study recently reported finding the juvenile of *P. parva* (20 mm standard length) in the estuary of the Volga River, although without morphological or genetic confirmation (Velikotskaya et al. 2025). Therefore, the colonisation of the Volga by this aggressive alien species might occur via different routes - along the Caspian Sea to Lower Volga (*P. parva* was found recently in the rivers of the Dagestan Region, close to the estuary of the Volga River – Barkhalov et al. (2023)) and via human-aided inter-basin fauna introductions in the Middle Volga / Don watershed. The alien species under consideration may rapidly expand their distribution within the Volga and Kama systems, as well as colonise the Baltic Sea Basin since the Volga serves as an invasive corridor from the Ponto-Caspian Basin to the Baltic Sea Basin (Dgebuadze 2011).

New natural (and/or introduced?) species

While the record of new alien species is anticipated in the Caspian Sea Basin (Dgebuadze 2011), the uncovering of a significant portion of novel natural diversity in the well-studied Volga Basin is surprising. This deserves consideration of each newly-recorded species for better understanding of the origin of this diversity, whether we overlooked natural diversity or detected new species because of their range expansion.

Family Salmonidae

Genus *Salmo*

Apart from previously known resident populations of *S. caspius* that inhabit the Middle Volga and Kama (Marić et al. 2016; Red Book of Russian Federation 2021), we detected an Atlantic lineage of *S. trutta* according to Bernatchez (2001) widely distributed in the neighbouring Baltic and White Sea Basins (=*S. trutta* s. str.). Moreover, individuals of the *S. trutta* s. str. co-occurred with *S. caspius* in the Yamanyelga River in the Kama system (locality no. 84 in Fig. 1) and were different in colour pattern (Fig. 6). In this locality, eight samples were identified as *S. caspius*, while three samples were identified as *S. trutta* s. str. The genetic divergence between these two lineages is rather low – as between

Figure 6. Resident forms of *Salmo trutta* (**A**) and *Salmo caspius* (**B**) cohabited the Yamanyalga River in the Kama Basin. Photo by O. Artaev. Scale bar: 10 mm.

most of *Salmo* spp. (e.g. Segherloo et al. (2021); Levin et al. (2022)), but it is enough to delineate these species. It is unclear whether *S. trutta* s. str. is native to the Volga Basin or was introduced and should be considered an invasive species. Another question is whether these two species hybridise in the zone of their secondary contact. Further genetic studies using nuclear markers are needed to clarify this issue. Since all *Salmo* populations in the Volga are threatened, clarifying their taxonomy and genetic distinctiveness is essential.

Family Gobionidae

Genus *Gobio*

One individual of gudgeon caught in the upper reach of a small river in the Middle Volga bordering with the Don River Basin was identified as *G. caucasicus* having the same haplotype as gudgeons from the Don Basin. The Don's gudgeon is considered an independent species *G. brevicirris* (Fricke et al. 2025), but has no genetic difference from *G. caucasicus*, which was described earlier from the Caucasus. Therefore, we preliminarily consider *G. brevicirris* as a junior synonym of *G. caucasicus*, but thorough revision is needed. Remarkably, the *G. caucasicus* was detected in the same river (locality no. 150 on Fig. 1) where alien species *Pseudorasbora parva* was (see above), which we consider as the case of repeated human-aided inter-basin introduction.

Family Leuciscidae

Genus *Leucaspius*

The genus is monotypic. The *L. delineatus* was described from the Danube Basin in Austria. The Volga's samples are genetically distant to the Danubian samples (K2P distances are over 3%) and can be considered as separated species.

Moreover, two lineages within Volga's samples were revealed with K2P distance exceeding 2% (Suppl. material 4). Since both lineages co-habit throughout the Volga, the revealed cryptic divergence might mirror the past secondary contact as was already demonstrated for the *Rutilus rutilus* and *R. lacustris* within the Volga and Kama systems (see Levin et al. (2017); Artaev et al. (2021)). Further morphological and genetic studies are needed to revise the complexity of this genus.

Genus *Leuciscus*

Apart from three known species for the Volga (*L. aspius*, *L. idus* and *L. leuciscus*), a fourth species – the Siberian dace *L. baicalensis* – was recorded in the Chusovaya River, Kama Basin. The main range of this species is located in Siberian rivers drained to the Arctic Ocean. Although the Chusovaya River belongs to the Kama-Volga Basin, its upper reaches are located in Siberia, east of the main Ural ridge and sharing a watershed with the Ob Basin (locality no. 67 in Fig. 1). Its first finding in the Chusovaya River extends its range to the Volga-Kama system and further highlights the important role of the Chusovaya River as a waterway for fauna exchange between Europe and Siberia (Makhrov et al. 2021; Artaev et al. 2024).

Genus *Phoxinus*

For a long time, *P. phoxinus* was listed for the Volga Basin. Recent revision showed that *P. isetensis* inhabits the Volga-Kama system in place of *P. phoxinus* (Artaev et al. 2024). This study revealed a novel lineage in the upper reaches of the Oka Basin (locality no. 136 in Fig. 1), which is highly divergent from the *P. isetensis* (5.9%). This cryptic diversity needs further investigation. We consider *Phoxinus* sp. from the Oka Basin as a candidate species.

Family Cobitidae

Genus *Cobitis* Linnaeus, 1758

Two species of the genus *Cobitis* were known from the Volga – *C. taenia* and *C. melanoleuca* (Bogutskaya et al. 2013; Vasil'eva et al. 2020). In addition, this study revealed two new species for the Volga basin: i) *C. tanaitica* and ii) *C. faridpaki*.

Cobitis tanaitica is distributed in the basins of the Black Sea and Sea of Azov, but was found to be rather widely distributed in the Lower and Middle Volga, as well as Kama systems by this study (Suppl. material 1: table S1). We suggest that this species might colonise the Volga Basin during one of the pluvial post-glacial periods of Pleistocene-Holocene when connection between Volga and adjacent basins occurred. The routes of dispersal of the *C. tanaitica* are likely similar to those revealed for the *Eudontomyzon mariae* which colonised Middle Volga from the Don Basin and *Eudontomyzon* sp. 'Dnieper' colonised Middle Volga from the Dnieper Basin (Levin et al. 2016). Likely, *C. tanaitica* also inhabits the part of the Western Dvina Basin, Baltic Sea drainage (Vasil'ev and Vasil'eva 2022). The *C. tanaitica* is a species involved in the interspecific hybridisation with other *Cobitis* spp. and generation of the diploid–polyploid hybrid complexes and the successive polyploidisation with transitions from sexual to unisexual forms experienced by triploid *Cobitis* hybrids (Vasil'ev et al. 1989; Choleva et al. 2014).

Cobitis faridpaki is a species inhabiting the rivers draining to the southern Caspian Sea in Iran (Mousavi-Sabet et al. 2011). Our samples were identified as *C. faridpaki* via NCBI BLASTn having a similarity of 99.69% to the samples from the Iranian part of the Caspian Basin – the Babol Rud, Karaj River and Tajan River (MK506143; MK506158; KY646316). Our finding of this species in the estuary of the Volga River (locality no. 198 in Fig. 1) greatly expands the range of this species. We suggest two explanations for the record of this species far from its known range. First, the species might have been shipped with ballast waters from the southern Caspian Sea to the northern Caspian Sea and, therefore, it is an alien species in the Volga system. Alternatively, this species may naturally have a wider distribution around the Caspian Sea than previously recognised. Given recently described *C. amphilekta* Vasil'eva & Vasil'ev, 2012 is locally distributed in Azerbaijan parts of the Caspian Sea Basin and assumedly in Kazakhstan's part of Caspian Sea (Vasil'eva and Vasil'ev 2012), one may suggest that *C. amphilekta* is conspecific with *C. faridpaki*, which is broadly distributed around the Caspian Sea. No DNA sequences of *C. amphilekta* are yet available to test this hypothesis.

Family Gobiidae

Genus *Proterorhinus*

Apart from *Proterorinus nasalis*, widely distributed in basins of Caspian Sea and Sea of Azov (Vasil'eva et al. 2025), a new unrecognised lineage *Proterorhinus* sp. not matching any yet described valid species (*P. blennioides*, *P. semipellucidus*, *P. victori*, *P. marmoratus*, *P. semilunaris*) was discovered in the Volga Estuary (locality nos. 186, 198 in Fig. 1). Genetically, it is identical to candidate cryptic species *Proterorhinus* sp. recorded by Neilson and Stepien (2009b) in the Chernozemelskiy canal (system of Manych-Don, Sea of Azov Basin). The species status of this genetic lineage, as well as its evolutionary history and routes of dispersal to the Volga Basin are yet to be clarified.

Species with no barcoding gap

No barcoding gap between some species of the genera *Leuciscus* and *Rutilus* was detected. In particular, low K2P distance (0.9%) was revealed between the sister pair of *L. leuciscus* and *L. idus*. For this pair of species, hybridisation and mitochondrial replacement have been reported (Hänfling and Brandl 2000; Zangl et al. 2022). Low levels of genetic divergence might also result from the past hybridisation between ancestral lineages. The minute distance between *R. lacustris* and *R. caspicus* in mtDNA (0.2%) is not surprising since it has already been reported, based on cytochrome *b* sequences of mtDNA (Levin et al. 2017). The explanation for this is an ancient and recent introgressive hybridisation between *Rutilus* spp. (see genomic study of Levin et al. (2025)).

Deep intraspecific distance – cryptic diversity or traces of the past hybridisation?

Five species showed relatively high intraspecies genetic distance. In particular, the Black and Caspian Sea sprat *Clupeonella cultriventris*, an alien species in

the Volga Basin, is genetically very diverse (2.21%). It points out: i) the complex evolutionary history of the species and/or ii) several donors of invasion to the Volga Basin (see also Karabanov et al. (2025)).

The belica '*Leucaspius delineatus*' inhabiting the Volga Basin is greatly different from *L. delineatus* in the type basin, the Danube River (see above) and might be a separate species. However, the Volga hosts two divergent lineages of *Leucaspius* (2.38%), which are not well grouped geographically (Suppl. material 4). Most likely, the Volga Basin is a zone of secondary contact of previously isolated lineages like fishes of the *Coregonus* and *Rutilus* spp. (Levin et al. 2017; Borovikova and Artamonova 2018). Alternatively, they might be the cryptic species. Both hypotheses should be tested further in morphological and genetic studies.

The ide *Leuciscus idus* showed a rather large intraspecific distance (1.42%) that is over the interspecific distance when compared to sister species *L. leuciscus* (see above). The possible reason is the past hybridisation with *L. leuciscus* and secondary contact of half-isolated ancient lineages of *L. idus* sensu lato.

The northern minnow *Phoxinus isetensis* has a moderately large intraspecific distance (1.26%). This species has the widest range amongst European *Phoxinus* minnows and likely geographically subdivided populations or sub-lineages (Artaev et al. 2024).

The Koshewnikow's sculpin *Cottus koshewnikowi* in the Volga system has a rather large intraspecific distance (1.10%) that is geographically structured (Volga and Kama). The cryptic diversity within this species needs further investigation.

Biogeographical implications

Biogeography of the Volga's fishes requires a separate thorough study. In this paper, we highlight the most important biogeographical issues revealed by this study. First, our results confirm the numerous inter-basin contacts and fish fauna exchange with the adjacent basins postulated previously (e.g. Marić et al. (2014); Levin et al. (2016); Makhrov et al. (2021)). In particular, results of this study corroborate colonisation of the Volga system by species from the following riverine basins: i) Arctic Ocean drainage via Siberian Ob' Basin in the Kama-Chusovaya corridor – record of the Siberian dace *Leuciscus baicalensis*; ii) Black Sea and Sea of Azov via the Don and Dnieper watersheds – records of *Gobio caucasicus*, *Cobitis tanaitica*, *Phoxinus* sp. and likely *Proterorhinus* sp.; iii) Southern Caspian Basin – record of *Cobitis faridpaki*. Based on newly-obtained and published data (e.g. Levin et al. (2016, 2017); Makhrov et al. (2021); Artaev et al. (2024)) on multiple and multidirectional past colonisation of the Volga River by various species from adjacent riverine and brackish water systems, we consider the Volga a large-scale melting pot. Likely, the mixing of fish fauna also resulted in the rise of extended zones of secondary contacts as revealed for the *Rutilus* spp. (Levin et al. 2025), a permanent hybrid system of reticulate speciation with diploid-triploid-tetraploid in the genus *Cobitis* (Vasil'ev et al. 1989) along with hybridizatioin within the genus *Coregonus* (Borovikova and Artamonova 2018). We are sure that the upcoming studies will include new instances of interspecific hybridisation in the Volga River Basin.

Despite significant mixing of the fish fauna in the Volga Basin caused by frequent inter-basin interactions during Pleistocene glaciations and deglaciations (Levin et al. 2017; Makhrov et al. 2021; this study), some indications of

regional endemism were also found. In particular, the sculpins *Cottus* show a divergence of the Kama Basin populations from the Volga populations at a level supposedly sufficient for the delimitation of a new species because this group is weakly variable in *COI* (e.g. Sideleva and Zhidkov (2024)). Species of the genera *Barbatula*, *Phoxinus* and *Leucaspius* also demonstrate some geographical structure. The first two shows that their Kama's populations are semi-isolated (recorded in Kama and the part of the Middle Volga that is close to the Kama). More studies involving larger sets of the samples need to clarify issues of local endemism or delimitation of geographically unique populations within the Volga and Kama riverine systems. Generally, high endemism of fish fauna in the Kama riverine system as revealed by this study is highlighted by the more ancient age of the Kama River (palaeo-Kama) compared to the Volga River (Sokolov 1952). Consequently, the Kama River system might serve as both a corridor for fish fauna exchange with Arctic Ocean drainage and as a refuge for endemic lineages during Pleistocene glaciations.

Conclusions

This study highlights the utility of barcoding for species identification of fish and lamprey species inhabiting the Volga River system. The presence of a barcoding gap for over 95% of analysed species confirmed the informativeness of the *COI* marker for species identification within the Volga Basin. Therefore, the generated *COI* sequences reference database can be successfully used in various disciplines – taxonomy, phylogeography, evolutionary history, biogeography, conservation genetics and food control along with studies using different types of DNA – museum DNA, ancient DNA in zooarchaeological studies. In particular, this study using the *COI* sequences has discovered seven new species for the Volga Basin not recorded previously and three novel evolutionary lineages, potentially new species. In addition, we showed divergent phylogeographic patterns reflecting different evolutionary history of species and highlighted the biogeography of the large water system in Eastern Europe. The generated *COI* dataset fills the large gap in the *COI* reference library of fishes inhabiting waterbodies of Eastern Europe and will largely contribute to the subsequent studies of European fish fauna. This dataset can also be used to amplify other mitochondrial genes, as other markers are also relevant for biodiversity monitoring, such as eDNA studies.

Acknowledgements

We are very grateful to Mikhail Borisov, Elena Borovikova, Yuri Gerasimov, Vladimir Gusakov, Oleg Ermakov, Dmitry Vekhov and Alexey Kutuzov for their help in sampling. We also acknowledge Arseniy Cherenkov for his help in the wet-lab.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Use of AI

No use of AI was reported.

Funding

This study was supported by the Russian Science Foundation, grant no. 24-44-20019 'Fishes of the Caspian Sea basin: genetic diversity, evolution and biogeography'.

Author contributions

Conceptualization: BAL. Data curation: BAL, MAL, ASK, NSM, AAB, AAG, IST, ONA. Formal analysis: AAG, ES, BAL. Funding acquisition: BAL. Investigation: MAL, KVL, AAG, BAL, SAP, ONA, NSM, ASK, AAB, ABR, IVP, IST, ES. Methodology: ES, ONA. Project administration: BAL, NSM, MAL. Resources: KVL, ASK, IST, SAP, ABR, IVP, MAL. Software: ES. Supervision: BAL. Validation: AAG, ES, MAL. Visualization: ES, ONA, AAG, BAL. Writing - original draft: BAL. Writing - review and editing: ES, IST, IVP, AAB, ABR, MAL, ASK, SAP, ONA, AAG, BAL, KVL, NSM.

Author ORCIDs

Boris A. Levin https://orcid.org/0000-0002-4044-2036
Aleksey A. Bolotovskiy https://orcid.org/0000-0003-2262-6808
Alexander A. Gandlin https://orcid.org/0000-0002-0125-1288
Nikolai S. Mugue https://orcid.org/0000-0001-8957-1931
Ivan V. Pozdeev https://orcid.org/0000-0002-3229-6972
Stepan A. Podolyako https://orcid.org/0000-0001-5357-5404

Data availability

All sequences were deposited to GenBank under Accession numbers PV840619–PV841769.

References

Aljanabi SM, Martinez I (1997) Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques. Nucleic Acids Research 25(22): 4692–4693. https://doi.org/10.1093/nar/25.22.4692

Appleyard SA, Ward RD, Pogonoski JJ, Graham A, Last PR, Deagle BE, Holmes B, Gomon MF, Bray DJ, Johnson JW, Hay AC, Moore GI, Hammer MP, Russell B, Graham KJ (2025) Australia's marine fishes DNA barcode reference library for integrated taxonomy, metabarcoding & eDNA research. Scientific Data 12(1): 21. https://doi.org/10.1038/s41597-025-04375-4

April J, Mayden RL, Hanner RH, Bernatchez L (2011) Genetic calibration of species diversity among North America's freshwater fishes. Proceedings of the National Academy of Sciences 108(26): 10602–10607. https://doi.org/10.1073/pnas.1016437108

Artaev ON, Ermakov OA, Vekhov DA, Konovalov AF, Levina MA, Pozdeev IV, Ruchin AB, Alyushin IV, Iljin VYu, Levin BA (2021) Genetic screening of distribution pattern of roaches *Rutilus rutilus* and *R. lacustris* (Cyprinidae) in broad range of secondary contact (Volga Basin). Inland Water Biology 14(2): 205–214. https://doi.org/10.1134/S1995082921020024

Artaev ON, Bolotovskiy AA, Turbanov IS, Gandlin AA, Kutuzov AV, Levina MA, Melentev DA, Pozdeev IV, Borisov MYa, Levin BA (2024) Forgotten for two centuries: redescription of *Phoxinus isetensis* (Georgi, 1775) (Cypriniformes, Leuciscidae) – the most

widespread minnow in Europe. Zoosystematics and Evolution 100(3): 1155–1173. https://doi.org/10.3897/zse.100.126702

Bandelt HJ, Forster P, Röhl A (1999) Median-joining networks for inferring intraspecific phylogenies. Molecular Biology and Evolution 16(1): 37–48. https://doi.org/10.1093/oxfordjournals.molbev.a026036

Barbarossa V, Bosmans J, Wanders N, King H, Bierkens MFP, Huijbregts MAJ, Schipper AM (2021) Threats of global warming to the world's freshwater fishes. Nature Communications 12: 1701. https://doi.org/10.1038/s41467-021-21655-w

Barkhalov RM, Stolbunov IA, Artaev ON, Turbanov IS, Rabazanov NI, Khlopkova MV, Karabanov DP (2023) Distribution of topmouth gudgeon *Pseudorasbora parva* (Cyprinidae) in waterbodies and watercourses of the Republic of Dagestan, Russia. Inland Water Biology 16(4): 781–787. https://doi.org/10.1134/S199508292304003X

Behrens-Chapuis S, Herder F, Esmaeili HR, Freyhof J, Hamidan NA, Özuluğ M, Šanda R, Geiger MF (2015) Adding nuclear rhodopsin data where mitochondrial COI indicates discrepancies–can this marker help to explain conflicts in cyprinids. DNA Barcodes 3(1): 187–199. https://doi.org/10.1515/dna-2015-0020

Behrens-Chapuis S, Herder F, Geiger MF (2021) Adding DNA barcoding to stream monitoring protocols – What's the additional value and congruence between morphological and molecular identification approaches? PLoS ONE 16(1): e0244598. https://doi.org/10.1371/journal.pone.0244598

Bengtsson J, Nilsson SG, Franc A, Menozzi P (2000) Biodiversity, disturbances, ecosystem function and management of European forests. Forest Ecology and Management 132(1): 39–50. https://doi.org/10.1016/S0378-1127(00)00378-9

Bernatchez L (2001) The evolutionary history of brown trout (*Salmo trutta* L.) inferred from phylogeographic, nested clade, and mismatch analyses of mitochondrial DNA variation. Evolution 55(2): 351–379. https://doi.org/10.1111/j.0014-3820.2001.tb01300.x

Bogutskaya NG, Kijaschko PV, Naseka AM, Orlova MI (2013) Identifications keys for fish and invertebrates. Vol. 1. Fish and molluscs. KMK Scientific Press Ltd., St. Petersburg-Moscow, 543 pp.

Borovikova EA (2017) Special traits of the genetic structure and origin of the population of vendace *Coregonus albula* of Pleshcheyevo Lake. Biology Bulletin 44(3): 245–250. https://doi.org/10.1134/S1062359017030037

Borovikova EA, Artamonova VS (2018) Morphological specificities of vendace (Salmoniformes: Salmonidae: Coregoninae: *Coregonus albula*) population in Lake Pleshcheyevo (the Volga River basin): relationships of two phylogenetic lineages in a new zone of secondary contact. Organisms Diversity & Evolution 18(3): 355–366. https://doi.org/10.1007/s13127-018-0375-5

Brown SDJ, Collins RA, Boyer S, Lefort M-C, Malumbres-Olarte J, Vink CJ, Cruickshank RH (2012) Spider: An R package for the analysis of species identity and evolution, with particular reference to DNA barcoding. Molecular Ecology Resources 12(3): 562–565. https://doi.org/10.1111/j.1755-0998.2011.03108.x

Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, Narwani A., Mace GM, Tilman D, Wardle DA,Kinzig AP, Daily GC, Loreau M, Grace JB, Larigauderie A, Srivastava DS, Naeem S. (2012) Biodiversity loss and its impact on humanity. Nature 486: 59–67. https://doi.org/10.1038/nature11148

Chen W, Ma X, Shen Y, Mao Y, He S (2015) The fish diversity in the upper reaches of the Salween River, Nujiang River, revealed by DNA barcoding. Scientific Reports 5(1): 17437. https://doi.org/10.1038/srep17437

Chernomor O, von Haeseler A, Minh BQ (2016) Terrace aware data structure for phylogenomic inference from supermatrices. Systematic Biology 65(6): 997–1008. https://doi.org/10.1093/sysbio/syw037

Choleva L, Musilova Z, Kohoutova-Sediva A, Paces J, Rab P, Janko K (2014) Distinguishing between incomplete lineage sorting and genomic introgressions: complete fixation of allospecific mitochondrial DNA in a sexually reproducing fish (*Cobitis*; Teleostei), despite clonal reproduction of hybrids. PLoS ONE 9(6): e80641. https://doi.org/10.1371/journal.pone.0080641

Dgebuadze YY (2011) A decade of research on invasions of alien species in the Holarctic. Russian Journal of Biological Invasions 2(2): 69. https://doi.org/10.1134/S207511171103012X

Dodds WK, Perkin JS, Gerken JE (2013) Human impact on freshwater ecosystem services: a global perspective. Environmental Science & Technology 47(16): 9061–9068. https://doi.org/10.1021/es4021052

Dong X, Ju T, Grenouillet G, Laffaille P, Lek S, Liu J (2020) Spatial pattern and determinants of global invasion risk of an invasive species, sharpbelly *Hemiculter leucisculus* (Basilesky, 1855). Science of the Total Environment 711: 134661. https://doi.org/10.1016/j.scitotenv.2019.134661

Dyldin YV, Hanel L, Fricke R, Orlov AM, Romanov VI, Plesnik, JAN, Interesova EA, Vorobiev DS, Kochetkova, M. O (2020) Fish diversity in freshwater and brackish water ecosystems of Russia and adjacent waters. Publications of the Seto Marine Biological Laboratory 45: 47–116. https://doi.org/10.5134/251251

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research 32(5): 1792–1797. https://doi.org/10.1093/nar/gkh340

Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek AR (1994) DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3(5): 294–299.

Fricke R, Eschmeyer WNR, van der Laan [Eds] (2025) Eschmeyer's catalog of fishes: Genera, species, references. (http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp). [Electronic version accessed 30 Sep 2025]

Froese R, Pauly D [Eds] (2025) FishBase. World Wide Web electronic publication. www.fishbase.org [Electronic version accessed 15 Sep 2025]

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Systematic Biology 59: 307–321. https://doi.org/10.1093/sysbio/syq010

Hänfling B, Brandl R (2000) Phylogenetics of European cyprinids: insights from allozymes. Journal of Fish Biology 57(2): 265–276. https://doi.org/10.1111/j.1095-8649.2000.tb02171.x

Haponski AE, Stepien CA (2013) Phylogenetic and biogeographical relationships of the *Sander* pikeperches (Percidae: Perciformes): patterns across North America and Eurasia. Biological Journal of the Linnean Society 110(1): 156–179. https://doi.org/10.1111/bij.12114

Hebert PD, Gregory TR (2005) The promise of DNA barcoding for taxonomy. Systematic Biology 54(5): 852–859. https://doi.org/10.1080/10635150500354886

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: Improving the ultrafast bootstrap approximation. Molecular Biology and Evolution 35(2): 518–522. https://doi.org/10.1093/molbev/msx281

Hooper DU, Chapin III FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM, Loreau M, Naeem S, Schmid B, Setälä H, Symstad AJ, Vandermeer J, Wardle D A (2005) Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. Ecological Monographs 75(1): 3–35. https://doi.org/10.1890/04-0922

Hubert N, Hanner R, Holm E, Mandrak NE, Taylor E, Burridge M, Watkinson D, Dumont P, Curry A, Bentzen P, Zhang J, April J, Bernatchez L (2008) Identifying Canadian freshwater fishes through DNA barcodes. PLoS ONE 3(6): e2490. https://doi.org/10.1371/journal.pone.0002490

IUCN (2025) Red List of Threatened Species. Version 2024-2. https://www.iucnredlist.org [Accessed on 30 Sep 2025]

Ivanova NV, Zemlak TS, Hanner RH, Hebert, PD (2007) Universal primer cocktails for fish DNA barcoding. Molecular Ecology Notes 7(4): 544–548. https://doi.org/10.1111/j.1471-8286.2007.01748.x

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods 14: 587–589. https://doi.org/10.1038/nmeth.4285

Karabanov DP, Kodukhova YV, Slyn'ko YV (2009) New findings of stone moroco *Pseudorasbora parva* (Temm. et Schl., 1846) in the European part of Russia. Russian Journal of Biological Invasions 1: 11–13. https://doi.org/10.1134/S2075111710030021

Karabanov DP, Bekker EI, Pavlov DD, Borovikova EA, Kodukhova YV, Kotov AA (2022) New sets of primers for DNA identification of non-indigenous fish species in the Volga-Kama basin (European Russia). Water 14(3): 437. https://doi.org/10.3390/w14030437

Karabanov DP, Pavlov DD, Dgebuadze YY, Bazarov MI, Borovikova EA, Gerasimov Y V, Kodukhova YV, Mikheev PB, Nikitin EV, Opaleva TL, Severov YA, Sabitova RZ, Smirnov AK, Solomatin YI, Stolbunov IA, Tsvetkov AI, Vlasenko SA, Voroshilova IS, Zhong W, Zhang X, Kotov AA (2023) A dataset of non-indigenous and native fish of the Volga and Kama Rivers (European Russia). Data 8(10): 154. https://doi.org/10.3390/data8100154

Karabanov DP, Kodukhova YV, Kotov AA (2025) Characteristics and Phylogenetic Significance of the Complete Mitochondrial Genome of *Clupeonella cultriventris* (Actinopterygii: Clupeiformes), a Nonindigenous Fish Species of the Rybinsk Reservoir (Volga River). Inland Water Biology 18(1): 220–226. https://doi.org/10.1134/S1995082924600972

Knebelsberger T, Dunz AR, Neumann D, Geiger MF (2015) Molecular diversity of Germany's freshwater fishes and lampreys assessed by DNA barcoding. Molecular Ecology Resources 15(3): 562–572. https://doi.org/10.1111/1755-0998.12322

Kodukhova YV, Karabanov DP (2021) Finding of Longtail Dwarf Goby *Knipowitschia longecaudata* (Actinopterygii: Gobiidae) in the Upstream Part of Unregulated Section of the Volga River. Inland Water Biology 14(5): 620–625. https://doi.org/10.1134/S1995082921050072

Kodukhova YV, Borovikova EA, Karabanov DP (2016) First record of stellate tadpole goby *Benthophilus stellatus* (Sauvage, 1874) (Actinopterygii: Gobiidae) in the Rybinsk Reservoir. Inland Water Biology 9(4): 428–430. https://doi.org/10.1134/S1995082916030081

Kottelat M, Freyhof J (2007) Handbook of European freshwater fishes (Vols 1–3). Cornol, Switzerland: Publications Kottelat, 646 pp.

Leigh JW, Bryant D (2015) POPART: full-feature software for haplotype network construction. Methods in Ecology & Evolution 6(9): 1110–1116. https://doi.org/10.1111/2041-210X.12410

Levin B, Ermakov A, Ermakov O, Levina M, Sarycheva O, Sarychev V (2016) Chapter Three. Ukrainian Brook Lamprey *Eudontomyzon mariae* (Berg): phylogenetic position, genetic diversity, distribution. Jawless Fishes of the World. Cambridge Scholars Publishing. Vol. 1, 58–82.

Levin BA, Simonov EP, Ermakov OA, Levina MA, Interesova EA, Kovalchuk OM, Malinina YA, Mamilov NS, Mustafayev NJ, Pilin DV, Pozdeev IV, Prostakov NI, Roubenyan HR, Titov SV, Vekhov DA (2017) Phylogeny and phylogeography of the roaches, genus *Rutilus* (Cyprinidae), at the Eastern part of its range as inferred from mtDNA analysis. Hydrobiologia 788(1): 33–46. https://doi.org/10.1007/s10750-016-2984-3

Levin BA, Simonov E, Matveyev MP, Artaev ON, Mustafayev NJ, Pashkov AN, Roubenyan HR (2018) DNA barcoding of the fishes of the genus *Alburnoides* (Actinopterygii, Cyprinidae) from Caucasus. Mitochondrial DNA Part A 29(1): 49–55. https://doi.org/10.1080/24701394.2016.1238900

Levin B, Simonov E, Gabrielyan BK, Mayden RL, Rastorguev SM, Roubenyan HR, Sharko FS, Nedoluzhko AV (2022) Caucasian treasure: Genomics sheds light on the evolution of half-extinct Sevan trout, *Salmo ischchan*, species flock. Molecular Phylogenetics and Evolution 167: 107346. https://doi.org/10.1016/j.ympev.2021.107346

Levin B, Simonov E, Šanda R, Vukić J, Zogaris S, Bolotovskiy A, Levina M, Mugue N, Mustafayev N, Merilä J (2025) Genomic Insights Into Diversity, Phylogeny, Hybridization and Evolutionary History of Palearctic Fish Genus *Rutilus* (Leuciscidae). Diversity and Distributions 31: e70124. https://doi.org/10.1111/ddi.70124

Makhrov AA, Vinarski MV, Gofarov MY, Dvoryankin GA, Novoselov AP, Bolotov IN (2021) Faunal exchanges between the basins of the Arctic Ocean and the Caspian Sea: Their history and current processes. Biology Bulletin 48(7): 892–906. https://doi.org/10.1134/S1062359021070190

Marić S, Askeyev IV, Askeyev OV, Monakhov SP, Bravničar J, Snoj A (2014) Phylogenetic and population genetic analysis of *Thymallus thymallus* (Actinopterygii, Salmonidae) from the middle Volga and upper Ural drainages. Hydrobiologia 740(1): 167–176. https://doi.org/10.1007/s10750-014-1951-0

Marić S, Askeyev O, Askeyev A, Monakhov S, Yanybaev N, Askeyev I, Galimova D, Snoj A (2016) Lack of mtDNA variation among remote middle Volga and upper Ural brown trout suggests recent and rapid recolonization. Journal of Applied Ichthyology 32(5): 948–953. https://doi.org/10.1111/jai.13126

Marques V, Castagné P, Polanco A, Borrero Pérez G H, Hocdé R, Guérin PE, Juhel JB, Velez L, Loiseau N, Letessier TB, Bessudo S, Valentini A, Dejean T, Mouillot D, Pellissier L, Villéger S (2021) Use of environmental DNA in assessment of fish functional and phylogenetic diversity. Conservation Biology 35(6): 1944–1956. https://doi.org/10.1111/cobi.13802

Meier R, Shiyang K, Vaidya G, Ng PK (2006) DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. Systematic Biology 55(5): 715–728. https://doi.org/10.1080/10635150600969864

Mendel J, Lusk S, Vasil'eva ED, Vasil'ev VP, Lusková V, Ekmekci FG, Erk'akan F, Ruchin A, Koščo J, Vetešník L, Halačka K, Šanda R, Pashkov AN, Reshetnikov SI (2008) Molecular phylogeny of the genus *Gobio* Cuvier, 1816 (Teleostei: Cyprinidae) and its contribution to taxonomy. Molecular Phylogenetics and Evolution 47(3): 1061–1075. https://doi.org/10.1016/j.ympev.2008.03.005

Meyer CP, Paulay G (2005) DNA barcoding: Error rates based on comprehensive sampling. PLoS Biology 3(12): e422. https://doi.org/10.1371/journal.pbio.0030422

Mineeva N, Lazareva V, Litvinov A, Stepanova I, Chuiko G, Papchenkov V, Korneva L, Scherbina G, Pryanichnikova E, Perova S, Gerasimov Y, Karabanov D, Levin B, Borovikova E, Voroshilova I, Tsvetkov A, Okhapkin A, Shurganova G, Dvinskikh S, Noskov V, Kitaev A, Alexevnina M, Istomina A, Presnova E, Seletkova E, Baklanov M, Zinov'ev E, Schletterer M (2022) The Volga River. In Rivers of Europe. Elsevier: Amsterdam, The Netherlands 27–79. https://doi.org/10.1016/B978-0-08-102612-0.00002-X

Minh, BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R (2020) IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. Molecular Biology and Evolution 37: 1530–1534. https://doi.org/10.1093/molbev/msaa015

Mousavi-Sabet H, Vasil'eva ED, Vatandoust S, Vasil'ev VP (2011) *Cobitis faridpaki* sp. nova – a new spined loach species (Cobitidae) from the Southern Caspian Sea basin (Iran). Journal of Ichthyology 51(10): 925–931. https://doi.org/10.1134/S0032945211100055

Mousavi-Sabet H, Ganjbakhsh B, Geiger MF, Freyhof J (2016) Redescription of *Gobio nigrescens* from the Hari River drainage (Teleostei: Cyprinidae). Zootaxa 4114(1): 71–80. https://doi.org/10.11646/zootaxa.4114.1.4

Muhala V, Guimarães-Costa A, Macate IE, Rabelo LP, Adam Rick Bessa-Silva AR, Watanabe L, dos Santos DG, Sambora L, Vallinoto M, Sampaio I (2024) DNA barcoding for the assessment of marine and coastal fish diversity from the Coast of Mozambique. PLoS ONE 19(2): e0293345. https://doi.org/10.1371/journal.pone.0293345

Mustafayev NJ, Ibrahimov SR, Levin BA (2015) Korean sharpbelly *Hemiculter leucisculus* (Basilewsky, 1855) (Cypriniformes, Cyprinidae) is a new species of Azerbaijan fauna. Russian Journal of Biological Invasions 6(4): 252–259. https://doi.org/10.1134/S2075111715040049

Neilson ME, Stepien CA (2009a) Escape from the Ponto-Caspian: Evolution and biogeography of an endemic goby species flock (Benthophilinae: Gobiidae: Teleostei). Molecular Phylogenetics and Evolution 52(1): 84–102. https://doi.org/10.1016/j.ympev.2008.12.023

Neilson ME, Stepien CA (2009b) Evolution and phylogeography of the tubenose goby genus *Proterorhinus* (Gobiidae: Teleostei): evidence for new cryptic species. Biological Journal of the Linnean Society 96(3): 664–684. https://doi.org/10.1111/j.1095-8312.2008.01135.x

Palandačić A, Kruckenhauser L, Ahnelt H, Mikschi E (2020) European minnows through time: museum collections aid genetic assessment of species introductions in freshwater fishes (Cyprinidae: *Phoxinus* species complex). Heredity 124(3): 410–422. https://doi.org/10.1038/s41437-019-0292-1

Panov VE, Dgebuadze YY, Shiganova TA, Filippov AA, Minchin D (2007) A risk assessment of biological invasions in the inland waterways of Europe: the Northern Invasion Corridor case study. In: Gherardi F (Eds) Biological invaders in inland waters: Profiles, distribution, and threats Springer Netherlands 639–656. https://doi.org/10.1007/978-1-4020-6029-8_35

Pereira AM, Levy A, Vukić J, Šanda R, Levin BA, Freyhof J, Geiger M, Choleva L, Francisco SM, Robalo JI (2021) Putting European lampreys into perspective: A global scale multilocus phylogeny with a proposal for a generic structure of the Petromyzontidae. Journal of Zoological Systematics and Evolutionary Research 59(8): 1982–1993. https://doi.org/10.1111/jzs.12522

Puillandre N, Lambert A, Brouillet S, Achaz G (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. Molecular Ecology 21: 1864–1877. https://doi.org/10.1111/j.1365-294X.2011.05239.x

Red Book of Russian Federation (2021) Vol. "Animals". 2nd edn. Moscow: "VNII Ecology" 1128 pp.

Ruban G, Khodorevskaya R, Shatunovskii M (2019) Factors influencing the natural reproduction decline in the beluga (*Huso huso*, Linnaeus, 1758), Russian sturgeon (*Acipenser gueldenstaedtii*, Brandt & Ratzeburg, 1833), and stellate sturgeon (*A. stellatus*, Pallas, 1771) of the Volga–Caspian basin: A review. Journal of Applied Ichthyology 35(1): 387–395. https://doi.org/10.1111/jai.13885

Sayer CA, Fernando E, Jimenez RR, Macfarlane NB, Rapacciuolo G, Böhm M, Brooks TM, Contreras-MacBeath T, Cox NA, Harrison I, Hoffmann M, Jenkins R, Smith KG, Vié JC, Abbott JC, Allen DJ, Allen GR, Barrios V, Boudot JP, Carrizo SF, Charvet P, Clausnitzer V, Congiu L, Crandall KA, Cumberlidge N, Cuttelod A, Dalton J, Daniels AG, De Grave S, De Knijf G, Dijkstra KDB, Dow RA, Freyhof J, García N, Gessner J, Getahun A, Gibson C, Gollock MJ, Grant MI, Groom AER, Hammer MP, Hammerson GA, Hilton-Taylor C, Hodgkinson L, Holland RA, Jabado RW Bignoli DJ, Kalkman VJ, Karimov BK, Kipping J, Kottelat M, Lalèyè PA,Larson HK, Lintermans M, Lozano F, Ludwig A, Lyons TJ, Máiz-Tomé L, Molur S, Ng HH, Numa C, Palmer-Newton AF, Pike C, Pippard HE, Polaz CNM, Pollock CM, Raghavan R, Rand PS, Ravelomanana T, Reis RE, Rigby CL, Scott JA, Skelton PH, Sloat MR, Snoeks J, Stiassny MLJ, Tan HH, Taniguchi Y, Thorstad EB, Tognelli MF, Torres AG, Torres Y, Tweddle D, Watanabe K, Westrip JRS, Wright EGE, Zhang E, Darwall WR (2025) One-quarter of freshwater fauna threatened with extinction. Nature 638: 138–145. https://doi.org/10.1038/s41586-024-08375-z

Schletterer M, Kuzovlev VV, Zhenikov YN, Tuhtan JA, Haidvogl G, Friedrich T, Górski K, Füreder L (2018) Fish fauna and fisheries of large European rivers: Examples from the Volga and the Danube. Hydrobiologia 814(1): 45–60. https://doi.org/10.1007/s10750-017-3370-5

Segherloo IH, Freyhof J, Berrebi P, Ferchaud AL, Geiger M, Laroche J, Levin BA, Normandeau E, Bernatchez L (2021) A genomic perspective on an old question: *Salmo* trouts or *Salmo trutta* (Teleostei: Salmonidae)? Molecular Phylogenetics and Evolution 162: 107204. https://doi.org/10.1016/j.ympev.2021.107204

Sheraliev B, Peng Z (2021) Molecular diversity of Uzbekistan's fishes assessed with DNA barcoding. Scientific Reports 11(1): 16894. https://doi.org/10.1038/s41598-021-96487-1

Sideleva VG, Zhidkov ZV (2024) Description of a new species of *Cottus* Linnaeus (Cottidae) from the Western Dvina/Daugava River system, Baltic Sea basin, based on integrative taxonomy. Zootaxa 5447(4): 473–495. https://doi.org/10.11646/zootaxa.5447.4.2

Slynko YV, Tereshchenko VG (2014) Freshwater fishes of the Ponto-Caspian Basin (diversity, faunogenesis, population dynamics, adaptation mechanisms). Polygraph-Plus, Moscow. https://doi.org/10.13140/2.1.1076.2560

Snyder MR, Stepien CA, Marshall NT, Scheppler HB, Black CL, Czajkowski KP (2020) Detecting aquatic invasive species in bait and pond stores with targeted environmental (e) DNA high-throughput sequencing metabarcode assays: Angler, retailer, and manager implications. Biological Conservation 245: 108430. https://doi.org/10.1016/j.biocon.2020.108430

Sokolov BS (1952) About the age of ancient sedimentary cover of the Russian platform. Bulletin of Academy of Sciences of USSR, Geology Series 5: 21–31. [in Russian]

Steinke D, Zemlak TS, Boutillier JA, Hebert PD (2009) DNA barcoding of Pacific Canada's fishes. Marine Biology 156: 2641–2647. https://doi.org/10.1007/s00227-009-1284-0

Tsoupas A, Papavasileiou S, Minoudi S, Gkagkavouzis K, Petriki O, Bobori D, Sapounidis A, Koutrakis E, Leonardos I, Karaiskou N, Triantafyllidis A (2022) DNA barcoding identification of Greek freshwater fishes. PLoS ONE 17(1): e0263118. https://doi.org/10.1371/journal.pone.0263118

Turbanov IS, Bolotovskiy AA, Artaev ON, Gandlin AA, Levina MA, Vasil'eva ED, Levin BA (2025) On the border of Europe and Asia: *Gobio uralensis*, a new species of gudgeons (Cypriniformes, Gobionidae) from the Caspian Sea basin. Zoosystematics and Evolution 101(2): 855–874. https://doi.org/10.3897/zse.101.147368

Ushio M, Murakami H, Masuda R, Sado T, Miya M, Sakurai S, Yamanaka H, Minamoto T, Kondoh, M. (2018) Quantitative monitoring of multispecies fish environmental DNA using high-throughput sequencing. Metabarcoding and Metagenomics 2: e23297. https://doi.org/10.3897/mbmg.2.23297

Vasil'ev VP, Vasil'eva KD, Osinov AG (1989) Evolution of a diploid-triploid-tetraploid complex in fishes of the genus *Cobitis* (Pisces, Cobitidae). Evolution and Ecology of Unisexual Vertebrates, Bulletin 466: 153–169.

Vasil'eva ED, Vasil'ev VP (2012) *Cobitis amphilekta* sp. nova, a new species of spined loaches (Cobitidae, Cypriniformes) from the Caspian Sea basin. Journal of Ichthyology 52(3): 200–206. https://doi.org/10.1134/S0032945212020154

Vasil'eva ED, Solovyeva EN, Levin BA, Vasil'ev VP (2020) *Cobitis derzhavini* sp. nova—a new spined loach species (Teleostei: Cobitidae) discovered in the Transcaucasia. Journal of Ichthyology 60(2): 135–153. https://doi.org/10.1134/S0032945220020198

Vasil'eva ED, Vasil'ev VP, Borovikova EA, Levina MA, Levin BA (2025) Species Diversity, Diagnostic Characters, and Distribution of Tubenose Gobies of the Genus *Proterorhinus* (Gobiidae): What are *P. semipellucidus* and *P. nasalis*? Journal of Ichthyology, 65: 737–752. https://doi.org/10.1134/S0032945225600508

Vasil'ev V, Vasil'eva E (2022) Triploid forms' karyotypes of spined loaches from the genus *Cobitis* (Actinopterygii: Cypriniformes: Cobitidae) of the upper Dnieper and Western Dvina rivers: Analysis of the triploids' origin. Acta Ichthyologica et Piscatoria 52: 67–75. https://doi.org/10.3897/aiep.52.81191

Velikotskaya PA, Podolyako SA, Litvinov KV (2025) The first find of the stone Moroko *Pseudorasbora parva* (Cyprinidae) in the Volga Delta. Russian Journal of Biological Invasions 16(1): 12–15. https://doi.org/10.1134/S2075111724700577

Wang T, Zhang YP, Yang ZY, Liu Z, Du YY (2020) DNA barcoding reveals cryptic diversity in the underestimated genus *Triplophysa* (Cypriniformes: Cobitidae, Nemacheilinae) from the northeastern Qinghai-Tibet Plateau. BMC Evolutionary Biology 20(1): 1–15. https://doi.org/10.1186/s12862-020-01718-0

Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PD (2005) DNA barcoding Australia's fish species. Philosophical Transactions of the Royal Society B: Biological Sciences 360(1462): 1847–1857. https://doi.org/10.1098/rstb.2005.1716

Ward RD, Hanner R, Hebert PD (2009) The campaign to DNA barcode all fishes, FISH BOL. Journal of Fish Biology 74(2): 329–356.https://doi.org/10.1111/j.1095-8649.2008.02080.x

Weiss SJ, Gonçalves DV, Secci-Petretto G, Englmaier GK, Gomes-Dos-Santos A, Denys GP, Persat H, Antonov A, Hahn C, Taylor EB, Froufe E (2021) Global systematic diversity, range distributions, conservation and taxonomic assessments of graylings (Te-

leostei: Salmonidae; *Thymallus* spp.). Organisms Diversity & Evolution 21(1): 25–42. https://doi.org/10.1007/s13127-020-00468-7

Xiong F, Shu L, Zeng H, Gan X, He S, Peng Z (2022) Methodology for fish biodiversity monitoring with environmental DNA metabarcoding: The primers, databases and bioinformatic pipelines. Water Biology and Security 1(1): 100007. https://doi.org/10.1016/j.watbs.2022.100007

Zangl L, Schäffer S, Daill D, Friedrich T, Gessl W, Mladinić M, Sturmbauer C, Wanzenböck J, Weiss SJ, Koblmüller S (2022) A comprehensive DNA barcode inventory of Austria's fish species. PLoS ONE 17(6): e0268694. https://doi.org/10.1371/journal.pone.0268694

Zardoya R, Doadrio I (1999) Molecular evidence on the evolutionary and biogeographical patterns of European cyprinids. Journal of Molecular Evolution 49(2): 227–237. https://doi.org/10.1007/PL00006545

Zemlak TS, Ward RD, Connell AD, Holmes BH, Hebert PD (2009) DNA barcoding reveals overlooked marine fishes. Molecular Ecology Resources 9: 237–242. https://doi.org/10.1111/j.1755-0998.2009.02649.x

Zhang Y, Qiu Y, Liu K, Zhong W, Yang J, Altermatt F, Zhang X (2024) Evaluating eDNA and eRNA metabarcoding for aquatic biodiversity assessment: From bacteria to vertebrates. Environmental Science and Ecotechnology 21: 100441. https://doi.org/10.1016/j.ese.2024.100441

Zhang Y, Zhang H, Akashi H, Albouy CP, Andres KJ, Barquín J, Brantschen J, Connon RE, Craine JM, Gleeson D, Goldenberg-Vilar A, González-Ferreras AM, Hatzenbuhler C, Hupało K, Hyde J, Iwasaki W, Johnson MD, Katz AD, Kuzovlev VV, Larson CE, Lecaudey LA, Leese F, Leray M, Li F, Macher T-H, Mauvisseau Q, Morán-Luis M, Nester G, Quintero H, Ravelomanana T, Chacko MR, Saccò M, Sales N, Schenekar T, Schletterer M, Schmidt S, Schulte NO, Schütz R, Sperry JH, Stevens ER, Stinson SA, Weiss S, Xia F, Zhang H, Zhang S, Zhong W, Zong S, Pellissier L, Zhang X, Altermatt F (2025) Integrated reanalysis of global riverine fish eDNA datasets shows robustness and congruence of biodiversity conclusions. bioRxiv, 2025-09. https://doi.org/10.1101/2025.09.16.676481

Supplementary material 1

Suplementary files

Authors: Boris A. Levin, Marina A. Levina, Oleg N. Artaev, Aleksey A. Bolotovskiy, Alexander A. Gandlin, Ilya S. Turbanov, Aleksandra S. Komarova, Nikolai S. Mugue, Ivan V. Pozdeev, Alexander B. Ruchin, Kirill V. Litvinov, Stepan A. Podolyako, Evgeniy Simonov

Data type: xlsx

Explanation note: Contains deveral Supplementary data on material studied, taxonomy, PCR conditions, data on intra- and intergenetic divergence, ASAP and ABGD partitions, images and haplotype network.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/mbmg.10.174947.suppl1

Supplementary material 2

Taxonomic pie chart

Authors: Boris A. Levin, Marina A. Levina, Oleg N. Artaev, Aleksey A. Bolotovskiy, Alexander A. Gandlin, Ilya S. Turbanov, Aleksandra S. Komarova, Nikolai S. Mugue, Ivan V. Pozdeev, Alexander B. Ruchin, Kirill V. Litvinov, Stepan A. Podolyako, Evgeniy Simonov

Data type: html

Explanation note: Interactive version of the plot in Fig. 2 'Pie chart showing the taxonomic composition of the examined COI dataset of fishes from the Volga River basin by family, genus, and species'.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/mbmg.10.174947.suppl2

Supplementary material 3

Supplementary image

Authors: Boris A. Levin, Marina A. Levina, Oleg N. Artaev, Aleksey A. Bolotovskiy, Alexander A. Gandlin, Ilya S. Turbanov, Aleksandra S. Komarova, Nikolai S. Mugue, Ivan V. Pozdeev, Alexander B. Ruchin, Kirill V. Litvinov, Stepan A. Podolyako, Evgeniy Simonov

Data type: jpg

Explanation note: Photographs of the new alien species revealed in the Volga basin - *Hemiculter leucisculus* and *Pseudorasbora parva*.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/mbmg.10.174947.suppl3

Supplementary material 4

Phylogeography

Authors: Boris A. Levin, Marina A. Levina, Oleg N. Artaev, Aleksey A. Bolotovskiy, Alexander A. Gandlin, Ilya S. Turbanov, Aleksandra S. Komarova, Nikolai S. Mugue, Ivan V. Pozdeev, Alexander B. Ruchin, Kirill V. Litvinov, Stepan A. Podolyako, Evgeniy Simonov

Data type: docx

Explanation note: Haplotype network of *Leucaspius* spp. inhabited the Volga basin and represented two lineages (K2P distance = 2.38%).

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/mbmg.10.174947.suppl4